



TITLE:

On the discrete Morse flow as a numerical tool (Variational Problems and Related Topics)

AUTHOR(S):

Omata, Seiro; Kinami, Shinichi

CITATION:

Omata, Seiro ...[et al]. On the discrete Morse flow as a numerical tool (Variational Problems and Related Topics). 数理解析研究所講究録 2001, 1181: 177-181

ISSUE DATE:

2001-01

URL:

<http://hdl.handle.net/2433/64561>

RIGHT:

On the discrete Morse flow as a numerical tool

Seiro OMATA and Shinichi KINAMI

小俣正朗 木南伸一

Department of Computational Science, Faculty of Science,
Kanazawa University
Kakuma-machi, Kanazawa 920-11 Japan

Dedicated to Professor Norio Kikuchi for his 60th birthday

1 Introduction

The **discrete Morse flow(DMF)** have been developed by Professor Norio Kikuchi(at Keio Univ.) for constructing solution of heat type equation related to a minimizing problem ([3], [4], [7], [8], [9], [10] and [11]). Kikuchi introduced the time-semidiscretized energy form below: For fixed positive constant h , consider the functional:

$$J_m(u) := \int_{\Omega} \frac{|u - u_{m-1}|^2}{h} dx + I(u), \quad (1.1)$$

for an elliptic functional $I(u)$. Here $\Omega(\subset \mathbf{R}^n)$ be a domain with Lipschitz boundary ($n \geq 1$). $I(u)$ is a functional which usually has the form

$$I(u) := \int_{\Omega} F(x, u, \nabla u) dx, \quad \text{on } \mathcal{K} := \{u : \Omega \rightarrow \mathbf{R}^N; u \in \text{suitable space}\} \quad (1.2)$$

where $N \geq 1$ and $F(x, z, p) : \Omega \times \mathbf{R}^N \times \mathbf{R}^{nN} \rightarrow \mathbf{R}$ with suitable regularity. Usually F is required to have good properties such as the elliptic structure for the first variation (See [2] for example.) We determine a sequence $\{u_m^h\}$ as minimizers of J_m in \mathcal{K} , called the **DMF**, inductively: Firstly, for an initial data $u_0 \in \mathcal{K}$ with $I(u_0) < \infty$, we define u_1 , as a minimizer of J_1 in \mathcal{K} . The next function $u_2 \in \mathcal{K}$ is determined a minimizer of J_2 in \mathcal{K} , and so on. $\{u_m^h\}$ is an approximate solution of a heat equation(See section 2) which possibly describes the Morse (semi)flow for $I(u)$.

Historically, minimizing method for heat equation was firstly introduced by Rektorys ([14]). Kikuchi rediscovered this type of method and introduced the form (1.1). By Kikuchi's rediscovery, many authors treated the problem of this type. For example, Bethuel et.al. treated harmonic mapping([1]), Nagasawa and Omata treated free boundary problem to seek a stationary point of $I(u)$ without passing the limit of $h \rightarrow 0$ ([12]). Koji Kikuchi constructed a varifold solution for non-convex functional $I(u)$. Hyperbolic type problems are also studied. (See [6] and [7].) Recently, Omata developed a numerical method using minimizing method (See [13] and its references.)

2 Convergence and asymptotic behavior

The important estimate on this flow is based on the following property:

$$J_m^h(u_m^h) \equiv \int_{\Omega} \frac{|u_m^h - u_{m-1}^h|^2}{h} dx + I(u_m^h) \leq J_m^h(u_{m-1}^h) \equiv I(u_{m-1}^h),$$

and therefore we have $\int_{\Omega} |u_m^h - u_{m-1}^h|^2 / h dx \leq I(u_{m-1}^h) - I(u_m^h)$. Summing up from $m = 1$ to M , we have a estimate:

$$I(u_M^h) + \sum_{m=1}^M \int_{\Omega} \frac{|u_m^h - u_{m-1}^h|^2}{h} dx \leq I(u_0). \quad (2.1)$$

This estimate is a basic estimate of this flow, from which main properties are obtained.

For showing a convergence theory, we define an approximate solution of a heat equation.

DEFINITION 2.1 We define functions \bar{u}^h and u^h on $\Omega \times (0, \infty)$ by

$$\bar{u}^h(x, t) = u_m^h(x), \quad u^h(x, t) = \frac{t - (m-1)h}{h} u_m^h(x) + \frac{mh - t}{h} u_{m-1}^h(x),$$

for $(x, t) \in \Omega \times ((m-1)h, mh]$.

Firstly, we mention convergence theory when h tends to zero. By use of (2.1), if F is **coercive** in $H^1(\Omega)$, we can easily obtain facts that the following norms are uniformly bounded with respect to h :

$$\begin{aligned} & \left\| \frac{\partial u^h}{\partial t} \right\|_{L^2((0, \infty) \times \Omega)}, \left\| \nabla \bar{u}^h \right\|_{L^\infty((0, \infty); L^2(\Omega))}, \left\| \nabla u^h \right\|_{L^\infty((0, \infty); L^2(\Omega))}, \\ & \left\| u^h \right\|_{L^\infty((0, \infty); L^2(\Omega))}, \left\| \bar{u}^h \right\|_{L^\infty((0, \infty); L^2(\Omega))}, \left\| u^h \right\|_{W^{1,2}((0, T) \times \Omega)}, \text{ (for all } T > 0). \end{aligned}$$

By use of these estimates, we see that by a suitable choice of subsequence $\{h_j\}$ ($h_j \rightarrow 0$, $j \rightarrow \infty$), approximate solutions may converge to a weak solution u (a limit function) in a some topology. Thus we can say:

THEOREM 2.2 If F satisfies good conditions, then a limit function is a weak solution, i.e.

$$\begin{aligned} \int_{\Omega} u_0 \eta(x, 0) dx &= \int_0^T \int_{\Omega} D_t u \eta dx dt \\ &+ \int_0^T \int_{\Omega} (F_{p_\alpha^i}(x, u, \nabla u) D_\alpha \eta^i + F_{z^i}(x, u, \nabla u) \eta^i) dx dt \end{aligned} \quad (2.2)$$

for all $\eta \in \overset{\circ}{W}_2^{1,1}((0, T) \times \Omega)$ with $\eta(x, T) = 0$, where $\overset{\circ}{V}_2((0, T) \times \Omega) = \{u \in L^2(Q_T), u_x \in L^2(Q_T); |u|_{Q_T} = \text{ess sup}_{0 \leq t \leq T} \|u(x, t)\|_{L^2(\Omega)} + \|u_x\|_{L^2(Q_T)} < \infty\}$.

For above theorem, very strong conditions for F is requested, so some mathematicians are trying to relax them. (See [9] for example.)

Secondly, we can get a result on asymptotic behavior of the D.M.F. $\{u_m\}$ as $m \rightarrow \infty$. From (2.1), we easily have $\|u_m - u_{m-1}\|_{L^2(\Omega)} \rightarrow 0$ as $m \rightarrow \infty$. Again, by (2.1), for any subsequence $\{u_{m_j}\} \subset \{u_m\}$, there exists a subsequence $\{u_{m_{j_\nu}}\} \subset \{u_{m_j}\}$ and a function u_∞ on Ω such that $u_{m_{j_\nu}} \rightarrow u_\infty$ weakly in $W^{1,2}(\Omega)$, and strongly in $L^2(\Omega)$, as $\nu \rightarrow \infty$. Moreover, we have:

THEOREM 2.3 If F satisfies good conditions, the limit function u_∞ is a minimizer of the functional

$$J_\infty(u) = \int_{\Omega} \left(\frac{|u - u_\infty|^2}{h} + F(x, u, \nabla u) \right) dx$$

in \mathcal{K} , hence, u_∞ satisfies $-\int_{\Omega} (F_{p_\alpha^i}(x, u, \nabla u) D_\alpha \phi^i + F_{z^i}(x, u, \nabla u) \phi^i) dx = 0$ for any $\phi \in C_0^\infty(\Omega)$.

Sometimes regularity for u is obtained from this minimality ([12]). In other word, u_∞ is not a simple stationary point to $I(u)$, but expected to have some regular properties.

3 Numerical Examples


By use of **DMF**, we can construct a simple algorithm (i.e. Nonlinear Optimization) for solving heat type equations numerically ([13]). Compared with the Galerkin method, in some problem, it works well and usually the structure would be simpler. We show two types of examples.

3.1 Ginzburg-Landau energy


In this experiment, we treat $(u : B_1(0) \rightarrow \mathbf{R}^2)$

$$I(u) = \int_{B_1(0)} \left(|\nabla u|^2 + \frac{1}{\delta} (|u|^2 - 1) \right) dx.$$

The boundary data is Dirichlet with 4 degrees and as a initial data we put 1 vortex with 4 degrees. The vortex splits into 4 vortices with 1 degree.



In this case, the boundary data is Neumann, and put 2 vortices with 1 degree as a initial data. They push each other and eventually they are going out.




3.2 A Free Boundary Problem

In these experiment, we treat $(u : B_1(0) \rightarrow \mathbf{R})$ the following free boundary $(\partial\{u > 0\})$ problem


$$I(u) = \int_{B_1(0)} (|\nabla u|^2 + \chi_{u>0}) dx.$$

The difference of boundary data causes a different conclusions.

In this experiment boundary data is high enough and the limit function becomes constant.



In this experiment the boundary data is low so that the limit function keeps the free boundary.



Acknowledgment

The author is grateful to professor Takashi Suzuki for giving us a chance to write paper in this Research Report in RIMS.

References

- [1] F. Bethuel - J. -M. Coron - J. -M. Ghidaglia - A. Soyeur, "Heat flows and relaxed energies for harmonic maps, in Nonlinear diffusion equations and their equilibrium states, 3. (Progress

- in nonlinear differential equations and their applications, 7.) Birkhäuser, Boston, Basel, Berlin (1992), 99-109.
- [2] Giaquinta, M., "*Multiple Integrals in the Calculus of Variations and Nonlinear Elliptic Systems*", Annals of Mathematical Studies, Princeton University Press.
 - [3] Haga, J., Kikuchi, N., "*On the higher integrability for the gradients of the solutions to difference partial differential systems of elliptic-parabolic type*", Z. Angew. Math. Phys., **51**, No.2, (2000), 290-303.
 - [4] Haga, J., Kikuchi, N., "*On the existence of the harmonic variational flow subject to the two-sided conditions*", Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), **243**, (1997), Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funktsii. 28. 324-337, 343.
 - [5] Hoshino, K., Kikuchi, N., "*Reverse Hölder inequality for functions with discrete-time variable*", Dvnam. Contin. Discrete Impuls. Systems, **6**, No.4, (1999), 503-517.
 - [6] Hoshino, K., Kikuchi, N., "*Gehring theory for time-discrete hyperbolic differential equations*", Comment. Math. Univ. Carolin., **39**, No.4, (1998), 697-707.
 - [7] Hoshino, K., Kikuchi, N., "*On a construction of weak solutions to linear hyperbolic partial differential systems with the higher integrable gradients*", Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), **233**, (1996), Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 27. 30-52, 255; translation in J. Math. Sci. (New York), **93**, No.5, (1999), 636-652.
 - [8] Ivanov. A. V., Kikuchi, N., Fraska. M., "*Partial regularity for quasilinear nonuniformly parabolic systems Some questions of mathematical physics and function theory*", J. Math. Sci. (New York), **92**, No.6, (1998), 4286-4301.
 - [9] Kikuchi, K., "*Constructing Weak Solutions in a Direct Variational Method and an Application of Varifold Theory*", J.D.E. **150**, 1-23 (1998).
 - [10] Kikuchi, N., "*An approach to the construction of Morse flows for variational functionals*", in "Nematics – Mathematical and Physical Aspects", ed: J. -M. Coron, J. -M. Ghidaglia, F. Hélein, NATO Adv. Sci. Inst. Ser. C: Math. Phys. Sci. **332**, Kluwer Acad. Publ., Dordrecht - Boston - London (1991), 195-198.
 - [11] Kikuchi, N., "*A method of constructing Morse flows to variational functionals*", Nonlinear World, **1** (1994), 131-147.
 - [12] Nagasawa, T., Omata, S., "*Discrete Morse semiflows of a functional with free boundary*", Adv. Math. Sci. Appl. **2** (1993), 147-187.
 - [13] Omata, S., Okamura, T., Nakane, K., "*Numerical analysis for the discrete Morse semiflow related to the Ginzburg Landau functional*", Nonlinear Analysis, **37**, No.5, (1999) 589-602.
 - [14] Rektorys, K., "*On application of direct variational methods to the solution of parabolic boundary value problems of arbitrary order in the space variables*", Czech. Math. J. **21**(96), (1971), 318-339.